

National Institute on Drug Abuse (NIDA) Alcohol Facts



<https://www.drugabuse.gov/>

Table of Contents

- [Alcohol Facts](#)
 - [Signs of Alcohol Misuse](#)
 - [Effects of Alcohol on Brains and Bodies](#)

Alcohol Facts

Drinks like beer, malt liquor, wine, and hard liquor contain alcohol. Alcohol is the ingredient that gets you drunk.

Hard liquor—such as whiskey, rum, or gin—has more alcohol in it than beer, malt liquor, or wine.

These drink sizes have about the same amount of alcohol in them:

- 12 ounces of regular beer
- 8-9 ounces of malt liquor
- 5 ounces of wine
- 1 ½ ounces of hard liquor



Being drunk can make a person feel very silly, angry, or sad for no reason. It can make it hard to walk in a straight line, talk clearly, or drive.

Photo by: [The National Institute of Alcohol Abuse and Alcoholism](#)

Some slang names for alcohol are:

- Booze
- Juice
- Hooch
- Sauce

NEXT: [Signs of Alcohol Misuse](#)

Signs of Alcohol Misuse



Photo by [©istock.com/KatarzynaBialasiewicz](https://www.istock.com/KatarzynaBialasiewicz)

Drinking alcohol is not necessarily a problem, but drinking too much can cause bad things to happen. People who drink too much alcohol might forget things that happened when they were drunk. This is called a blackout.

People drunk on alcohol often:

- laugh and talk loudly
- feel dizzy
- have blurry vision
- have trouble staying on their feet and sway when they walk
- slur words when they talk
- feel sleepy and relaxed
- pass out
- throw up
- get violent

Getting drunk can lead you to do or say things that you regret later on. It also makes you more likely to get into a car crash and get hurt.

After drinking a lot, people get a headache and feel sick. This is called a hangover.

People who misuse alcohol, or people who are addicted, start having to drink more and more to get drunk. They might have a drink in the morning to calm down or stop a hangover. They might drink alone, and they might keep it a secret.

They might forget things that happened when they were drunk. This is called a blackout.

NEXT: [Effects of Alcohol on Brains and Bodies](#)

Effects of Alcohol on Brains and Bodies



Photo by ©istock.com/[aijohn784](#)

Being drunk makes you more likely to get hurt or killed. Alcohol is involved in many deadly car crashes.

These are just some of the problems alcohol can cause:

Diseases

Heavy drinking over the years can raise your risk for stroke (a “brain attack”), cancer, liver disease, and other illnesses.

People can forget to use condoms when they're drunk. They can then have unsafe sex and get HIV/AIDS or hepatitis (a liver disease).

Problems with Your Baby

If a pregnant woman drinks alcohol, it can cause brain damage, which can lead to other health problems in the baby.

Accidents or Death

Being drunk makes you more likely to get hurt or killed. Alcohol is involved in:

- 60 percent of drownings, murders, and deadly burns
- 50 percent of severe injuries and sexual attacks
- 40 percent of deadly driving crashes, deadly falls, and suicides

Brain Damage

Heavy drinking over the years can permanently hurt your brain cells. This can make it hard to walk, remember, or learn new things.

Alcohol Poisoning

You can die from drinking a lot of alcohol at one time, called alcohol poisoning. It can especially happen to young people who are not used to drinking.

Signs of alcohol poisoning:

- confusion
- throwing up
- slowed breathing
- cold, blue skin
- coma (you can't wake them up)
- seizures (when your body starts twitching and jerking uncontrollably)

If someone you know has any of these signs, you should call 911 right away.

Addiction

You can get addicted to alcohol just like other drugs. Over time, alcohol can change the way your brain works. If you stop drinking alcohol, your body can get confused and you can start to feel really sick. This makes it hard to stop. This is called addiction.

People who are trying to quit drinking might:

- feel nervous and sad
- shake
- sweat
- have trouble sleeping

They will feel a very strong need to drink alcohol.

Fortunately, there are medicines and other treatments that can help someone recover from alcohol addiction.

Remember that even if you get treatment, it can be hard to stop drinking. People who stopped using alcohol can still feel strong cravings to have a drink, sometimes even years later. This is normal. This is also why it's important to stay in treatment for as long as your doctor suggests, and

learn to enjoy friendships and activities that do not center on drinking.